

Detecting Cysteine Modifications in methanogen *Methanosarcina mazei* Gö1

Rachel Ogorzalek Loo^{1*} (rloo@mednet.ucla.edu), Phuong H.N. Nguyen^{1,2}, Hong Hanh Nguyen¹, **Todd Yeates**¹, Joseph Loo¹, and Robert Gunsalus¹

¹University of California-Los Angeles, Los Angeles, CA, USA; ²University of Science, VNU-HCMUS, Vietnam

<http://www.doe-mbi.ucla.edu/>

Project Goals: To elucidate the biological pathways of microbes relevant to microbial biofuel production and to global carbon cycling. These studies employ proteomics and mass spectrometry to characterize protein post-translational modifications.

Archaea in genus *Methanosarcina* are distributed broadly from marine to fresh water environments. They produce methane from a wide range of substrates including acetate, methylamines, and methanol and account for a large percentage of global methane emission. In methanogenesis, several important steps rely on thiol intermediates; *e.g.*, methyl transfer from tetrahydrosarcinopterin (H4SPT) to coenzyme M (mercaptoethanesulfonate), methane release by oxidation of coenzyme M and coenzyme B to form a heterodimer, and recycling of coenzymes M and B after reduction by heterodisulfide reductase. The importance of thiols to methanogenesis encouraged us to explore cysteine modifications in *Methanosarcina mazei*.

Tryptic peptides were generated with and without reduction/alkylation from cell lysates of *Methanosarcina* cultivated on methanol and on other carbon substrates. Peptides were analyzed by LC-MS/MS to identify proteins and to inventory post-translational modifications. Among the most abundant modifications observed was cysteinylation (Cys+119), identified on over 40 of proteins. Protein cysteinylation was observed not only from cultures maintaining reducing conditions with Na₂S/cysteine addition, but also from those supplementing with Na₂S only. Other modifications detected included Cys+30 (trisulfide in multi-cysteine peptides), Cys+140, Cys+151, and Cys+152. Modified cysteines appeared in active sites of some metabolic enzymes. The significance of these modifications is being explored.

This work was supported by the Department of Energy Office of Science (BER) through DE-FC02-02ER63421 to the UCLA-DOE Institute.