

Leveraging *Agave* and *Kalanchoë* Genomics Resources to Transfer Crassulacean Acid Metabolism (CAM) Modules into C₃ Species Using Synthetic Biology Approaches

Xiaohan Yang¹(yangx@ornl.gov), Rongbin Hu¹, Degao Liu¹, Henrique Cestari De Paoli, Paul E. Abraham², John Cushman³, Anne M. Borland^{1,4}, Gerald Tuskan¹

¹Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, TN;

²Chemical Sciences Division, Oak Ridge National Laboratory, Oak Ridge, TN

³Department of Biochemistry and Molecular Biology, University of Nevada, Reno, NV

⁴School of Biology, Newcastle University, Newcastle, UK

<http://cambiodesign.org/>

Project Goals: Crassulacean acid metabolism (CAM) is a specialized mode of photosynthesis that features a temporal CO₂ pump with nocturnal CO₂ uptake, facilitates increased water-use efficiency (WUE), and enables CAM plants to inhabit water-limited semi-arid or seasonally dry environments. CAM provides an excellent opportunity for engineering both enhanced WUE and photosynthetic performance into bioenergy crops. This project has two main goals: 1) to identify the CAM-associated genes and gene networks using systems biology approaches and 2) to engineer CAM gene modules into C₃ species using synthetic biology approaches. The success of the project could allow biomass production on semi-arid, abandoned, or marginal agricultural lands.

CAM-into-C₃ engineering requires multiple CAM-related genes to be manipulated in a modular manner, including: 1) a carboxylation module for CO₂ fixation and nocturnal accumulation of malic acid in the vacuole; 2) a decarboxylation module for release of CO₂ from malate; and 3) a stomatal control module for nocturnal stomatal opening and stomatal closure during the daytime (Yang et al. 2015). We have generated rich genomics resources for two important CAM models *Agave* (Abraham et al. 2016) and *Kalanchoë* (phytozome.jgi.doe.gov). Our comparative analysis of protein sequences and gene expression data have identified CAM-related genes in *Agave* and *Kalanchoë*. To characterize the function of CAM-related genes, the CRISPR/Cas9-based genome-editing technology (Liu et al. 2016) was used to create loss-of-function mutants in *K. fedtschenkoi*. Some of the genes related to carboxylation, decarboxylation, and stomatal movement in *Agave* and *Kalanchoë* have been engineered into three C₃ species as described below.

Engineering of carboxylation module: Genes related to carboxylation in *Agave* and *Kalanchoë*, including β -type carbonic anhydrase (β -CA), phosphoenolpyruvate carboxylase (PEPC), PEPC kinase (PPCK), malate dehydrogenase (MDH) and tonoplast aluminium-activated malate transporter (ALMT), were successfully introduced individually into C₃ species including model plants (i.e., *Arabidopsis* and tobacco) and bioenergy crop *Populus*. The transgenic plants were characterized using RT-PCR and western blot analysis to validate the expression of the transgenes. Phenotypical characterization of the transgenic plants is being conducted.

Engineering of decarboxylation module: Genes related to decarboxylation in *Agave* and *Kalanchoë*, including tonoplast dicarboxylate transporter (tDT), pyruvate phosphate dikinase (PPDK) and NADP-dependent malic enzyme (NADP-ME), were successfully introduced into C₃ species including model plants (i.e., *Arabidopsis* and tobacco) and the bioenergy crop *Populus*.

The transgenic plants were characterized using RT-PCR and western blot analysis to validate the expression of the transgenes. Phenotypical characterization of the transgenic plants is being conducted. Also, we made use of transgenic plants (*amiRs*), which show timely increase of malate by ≈ 2 -fold at early night, to reprogram decarboxylation using synthetic biology principles. First, we complemented *amiRs*, which are partially deficient in CO₂ release from malate, with a dark-induced promoter driving a functional NADP-ME enzyme and monitored gas exchange during a 20-hour period. Next, we segregated the functional NADP-ME out and re-evaluated CO₂ release from malate in the reverted line (*amiR-r*). Our results demonstrate the robustness of a controller to effectively link 3 inputs, which includes an ON-OFF switch, and generate two distinct outputs for diel control of intracellular CO₂ in leaves. This capability has different applications including the making of future CAM-engineered plants inducible under a specific input (e.g., drought).

Engineering of stomatal control module: Two strategies were used to change the stomatal behavior of C₃ plants: 1) transfer of multiple stomatal movement-associated genes from *K. fedtschenkoi* to C₃ plants and 2) synthesizing AND gate genetic circuits based upon the CRISPR-dCas9 system. More than 10 genetic circuits targeting ‘CAM-like stomatal movement’ were designed, assembled, and transferred into three target C₃ species (i.e., *Arabidopsis*, tobacco, and poplar) via *Agrobacterium*-mediated transformation. Transgenic plants were generated, and identified by PCR, qRT-PCR, and western blot analysis. The effects of these genetic circuits on stomatal movement in the transgenic plants will provide useful information for optimizing the strategy of CAM-into-C₃ engineering. Also, success of this research could facilitate future efforts to engineer other multi-gene traits through synthetic biology approaches.

This material is based upon work supported by the Department of Energy, Office of Science, Genomic Science Program (under award number DE-SC0008834). Oak Ridge National Laboratory is managed by UT-Battelle, LLC for the US Department of Energy (under contract number DE-AC05-00OR22725).

References

- Abraham PE, Yin H, Borland AM, Weighill D, Lim SD, De Paoli HC, Engle N, Jones PC, Agh R, Weston DJ et al. 2016. Transcript, protein and metabolite temporal dynamics in the CAM plant Agave. *Nature Plants* **2**: 16178.
- Liu D, Hu R, Palla KJ, Tuskan GA, Yang X. 2016. Advances and perspectives on the use of CRISPR/Cas9 systems in plant genomics research. *Curr Opin Plant Biol* **30**: 70-77.
- Yang X, Cushman JC, Borland AM, Edwards EJ, Wullschleger SD, Tuskan GA, Owen NA, Griffiths H, Smith JAC, De Paoli HC et al. 2015. A roadmap for research on crassulacean acid metabolism (CAM) to enhance sustainable food and bioenergy production in a hotter, drier world. *New Phytologist* **207**: 491-504.